

From Flows and Metrics to Dynamics

C.Udriște and A.Udriște

University Politehnica of Bucharest
 Department of Mathematics I
 Splaiul Independenței 313
 77206 Bucharest, Romania
 email:udriste@mathem.pub.ro

Abstract

Recall that a vector field on an n -dimensional differentiable manifold M is a mapping X defined on M with values in the tangent bundle TM that assigns to each point $x \in M$ a vector $X(x)$ in the tangent space T_xM . A vector field may be interpreted alternatively as the right-hand side of an autonomous system of first-order ordinary differential equations, i.e., a flow. Now we show that any flow can be enveloped by a conservative dynamics using a semi-Riemann metric g on M . This kind of dynamics was called *geometric dynamics* [7]-[9].

The given vector field, the initial semi-Riemann metric, the Levi-Civita connection, and an associated (1,1)-tensor field are used to build a new geometric structure (e.g., semi-Riemann-Jacobi, semi-Riemann-Jacobi-Lagrange, semi-Finsler-Jacobi, etc) on the manifold M ensuring that all the trajectories of a geometric dynamics are pre-geodesics (Lorentz-Udriște world-force law). Implicitly, we solved a problem rised first by Poincaré: find a suitable geometric structure that converts the trajectories of a given vector field into geodesics.

Mathematics Subject Classification 2000: 70G45, 70S05, 53C50

Key words: flows, metrics, geometric dynamics, Hamiltonian equations

1 Geometric Dynamics

Let M be an n -dimensional differentiable manifold. A C^∞ vector field X on M defines the flow

$$(1) \quad \frac{dx}{dt} = X(x).$$

A semi-Riemann metric g on the manifold M is a C^∞ symmetric tensor field of type (0,2) which assigns to each point $x \in M$ a nondegenerate inner product $g(x)$ on the tangent space $T_x M$ of signature (r, s) . The pair (M, g) is called a *semi-Riemann manifold*.

The vector field X and the semi-Riemann metric g determine the *energy*

$$f : M \rightarrow R, \quad f = \frac{1}{2}g(X, X).$$

The vector field (flow) X on (M, g) is called:

- 1) *timelike*, if $f < 0$;
- 2) *nonspacelike or causal*, if $f \leq 0$;
- 3) *null or lightlike*, if $f = 0$;
- 4) *spacelike*, if $f > 0$.

Let X be a nowhere vanishing vector field, of everywhere constant energy. Upon rescaling, it may be supposed that $f \in \{-1, 0, 1\}$. Generally, $\mathcal{E}$ is the set of zeros of the vector field X . If $\mathcal{E} \neq \emptyset$, then this rescaling is possible only on $M \setminus \mathcal{E}$.

Let ∇ be the Levi-Civita connection of (M, g) . Using the operator $\frac{\nabla}{dt}$ (covariant differentiation along a solution) we obtain the prolongation

$$(2) \quad \frac{\nabla}{dt} \frac{dx}{dt} = \nabla_{\frac{dx}{dt}} X$$

of the differential system (1) or of any perturbation of the system (1) obtained adding to the second member X a parallel vector field Y with respect to the covariant derivative ∇ . The prolongation by derivation represents the general dynamics of the flow. The vector field Y can be used to illustrate a progression from stable to unstable flows, or converse.

The vector field X , the metric g , and the connection ∇ determine the external (1,1)-tensor field

$$F = \nabla X - g^{-1} \otimes g(\nabla X),$$

$$F_j^i = \nabla_j X^i - g^{ih} g_{kj} \nabla_h X^k, \quad i, j, h, k = 1, \dots, n,$$

which characterizes the *helicity* of vector field (flow) X .

First we write the differential system (2) in the equivalent form

$$(2') \quad \frac{\nabla}{dt} \frac{dx}{dt} = g^{-1} \otimes g(\nabla X) \left(\frac{dx}{dt} \right) + F \left(\frac{dx}{dt} \right).$$

Successively we modify the differential system (2') as follows:

$$(3) \quad \frac{\nabla}{dt} \frac{dx}{dt} = g^{-1} \otimes g(\nabla X)(X) + F \left(\frac{dx}{dt} \right),$$

$$(4) \quad \frac{\nabla}{dt} \frac{dx}{dt} = g^{-1} \otimes g(\nabla X) \left(\frac{dx}{dt} \right) + F(X),$$

$$(5) \quad \frac{\nabla}{dt} \frac{dx}{dt} = g^{-1} \otimes g(\nabla X)(X) + F(X).$$

Obviously, the second order systems (3), (4), (5) are prolongations of the first order system (1). Each of them is connected either to the dynamics of the field X or to the dynamics of a particle which is sensitive to the vector field X . Since

$$g^{-1} \otimes g(\nabla X)(X) = \text{grad } f,$$

we shall show that the prolongation (3) describes a conservative dynamics of the vector field X or of a particle which is sensitive to the vector field X . The physical phenomenon produced by (4) or (5) was not yet studied.

Theorem. 1) If $F = 0$, then the kinematic system (1) prolonges to a potential dynamical system with n degrees of freedom, namely

$$(3') \quad \frac{\nabla}{dt} \frac{dx}{dt} = \text{grad } f.$$

2) If $F \neq 0$, then the kinematic system (1) prolonges to a non-potential dynamical system with n degrees of freedom, namely

$$(3'') \quad \frac{\nabla}{dt} \frac{dx}{dt} = \text{grad } f + F \left(\frac{dx}{dt} \right).$$

Corollary. *If the metric g is chosen such that $f \in \{-1, 0, 1\}$ on $M \setminus \mathcal{E}$, then the flow generated by X is global and the dynamical systems $(3')$, $(3'')$ are reduced to*

$$\frac{\nabla^2 x}{dt^2} = 0, \quad \frac{\nabla^2 x}{dt^2} = F\left(\frac{dx}{dt}\right).$$

Let us show that the dynamical systems $(3')$ and $(3'')$ are conservative. To simplify the exposition we identify the tangent bundle TM with the cotangent bundle T^*M using the semi-Riemann metric g .

Theorem. 1) *The trajectories of the dynamical system $(3')$ are the extremals of the Lagrangian*

$$L = \frac{1}{2}g\left(\frac{dx}{dt}, \frac{dx}{dt}\right) + f(x).$$

2) *The trajectories of the dynamical system $(3'')$ are the extremals of the Lagrangian*

$$L = \frac{1}{2}g\left(\frac{dx}{dt} - X, \frac{dx}{dt} - X\right) = \frac{1}{2}g\left(\frac{dx}{dt}, \frac{dx}{dt}\right) - g\left(X, \frac{dx}{dt}\right) + f(x).$$

3) *The dynamical systems $(3')$ and $(3'')$ are conservative, the Hamiltonian being the same for both cases, namely*

$$H = \frac{1}{2}g\left(\frac{dx}{dt}, \frac{dx}{dt}\right) - f(x).$$

The restriction of the Hamiltonian H to the flow of the vector field X is zero. Obviously the values of the Hamiltonian H can be positive, negative or zero, even if the metric g is a Riemannian metric; therefore, just in this case, there exist boundary-value problems associated to the differential system (3), having three solutions (for example, the first corresponding to constant total energy $H < 0$, the second for $H = 0$, and the third for $H > 0$).

For the next theorem we recall that a *pregeodesic* is a smooth curve which may be reparametrized to be a geodesic.

Theorem (Lorentz-Udriște World-Force Law). 1) *Every non-constant trajectory of the dynamical system $(3')$, which corresponds to a constant value H_0 of the Hamiltonian, is a pregeodesic of the semi-Riemann-Jacobi manifold*

$$(M \setminus \mathcal{E}, \overleftarrow{\mathcal{F}} = (\mathcal{H}, + \{ \})).$$

2) Let g_{ij} be the local components of the metric g and let Γ_{jk}^i , $i, j, k = 1, \dots, n$ be the local components of the connection ∇ . Every non-constant trajectory of the dynamical system (3''), which corresponds to a constant value H_0 of the Hamiltonian, is a horizontal pregeodesic of the semi-Riemann-Jacobi-Lagrange manifold

$$(M \setminus \mathcal{E}, \mathcal{F} = (\mathcal{H}_l + \{\})\}, \mathcal{N}_l^\rangle = -\rangle_{||}^\dagger - \mathcal{F}_l^\rangle, \quad \rangle, |, || = \infty, \dots, \setminus).$$

Corollary. If the metric g is chosen such that $f \in \{-1, 0, 1\}$ on $M \setminus \mathcal{E}$ and if we denote

$$\alpha^2 = g\left(\frac{dx}{dt}, \frac{dx}{dt}\right), \quad \beta = k^{1/2} g\left(X, \frac{dx}{dt}\right),$$

then every trajectory of the dynamical system (3''), with $\alpha^2 k^{1/2} + \beta = 0$, is a pregeodesic of the semi-Finsler-Jacobi manifold

$$(M \setminus \mathcal{E}, \mathcal{L} = \mathcal{F}_{\Xi}^\Xi = \Xi \alpha \beta, \quad \Xi = \text{constant}).$$

2 Hamiltonian Structures on the Tangent Bundle

Let N be a $2n$ -dimensional manifold. A nondegenerate and closed 2-form $\wedge$ on N is called *symplectic form*. A manifold N with a given symplectic form is called a *phase space*.

Let $(N, \wedge)$ be a phase space, and $H : N \rightarrow \mathbb{R}$ be a C^∞ real function. We define the Hamilton gradient X_H as being the vector field which satisfies

$$\wedge_p(X_H(x), v) = dH(x)(v), \quad \forall v \in T_x N,$$

and the Hamilton equations as

$$\frac{dx}{dt} = X_H(x).$$

Let (M, g) be a semi-Riemann manifold with n dimensions. Let X be a C^∞ vector field on M , and $\omega = g \circ F$ the 2-form associated to the tensor field $F = \nabla X - g^{-1} \otimes g(\nabla X)$ via the metric g .

The tangent bundle is usually endowed with the Sasaki metric G created by g . If (x^i, y^i) are the coordinates of the point $(x, y) \in TM$ and Γ_{jk}^i are the components of the connection induced by g_{ij} , then

$$\left(\frac{\partial}{\partial x^i} - \Gamma_{ij}^h y^j \frac{\partial}{\partial y^h}, \frac{\partial}{\partial y^i} \right), \quad (dx^j, \delta y^j = dy^j + \Gamma_{hk}^j y^k dx^h)$$

are dual frames. Also the metric of Sasaki transcribes

$$G = g_{ij} dx^i \otimes dx^j + g_{ij} \delta y^i \otimes \delta y^j.$$

Theorem. *The dynamical system $(3')$ lifts to TM as a Hamilton dynamical system with respect to the Hamiltonian*

$$H = \frac{1}{2}g \left(\frac{dx}{dt}, \frac{dx}{dt} \right) - f(x)$$

and the symplectic 2-form

$$\Omega_1 = g_{ij} dx^i \wedge \delta y^j.$$

Hint. $\eta_1 = g_{ij} y^i dx^j$, and $d\eta_1 = -\Omega_1$.

Theorem. *The dynamical system $(3'')$ lifts to TM as a Hamilton dynamical system with respect to the Hamiltonian*

$$H = \frac{1}{2}g \left(\frac{dx}{dt}, \frac{dx}{dt} \right) - f(x)$$

and the symplectic 2-form

$$\Omega_2 = \frac{1}{2} \omega_{ij} dx^i \wedge dx^j + g_{ij} dx^i \wedge \delta y^j.$$

Hint. $\eta_2 = -g_{ij} X^i dx^j + g_{ij} y^i dx^j$, and $d\eta_2 = -\Omega_2$.

Pendulum Geometric Dynamics. We use the Riemannian manifold (R^2, δ_{ij}) . The small oscillations of a plane pendulum are described as solutions of the differential system (plane pendulum flow)

$$(6) \quad \frac{dx_1}{dt} = -x_2, \quad \frac{dx_2}{dt} = x_1.$$

In this case $x_1(t) = 0$, $x_2(t) = 0$, $t \in R$ is the equilibrium point and $x_1(t) = c_1 \cos t + c_2 \sin t$, $x_2(t) = c_1 \sin t - c_2 \cos t$, $t \in R$ is the general solution (family of circles with same centre).

Let

$$X = (X_1, X_2), \quad X_1(x_1, x_2) = -x_2, \quad X_2(x_1, x_2) = x_1,$$

$$f(x_1, x_2) = \frac{1}{2}(x_1^2 + x_2^2), \quad \text{rot } X = (0, 0, 2), \quad \text{div } X = 0.$$

The pendulum flow conserves the areas. The prolongation by derivation of the kinematic system (6) is

$$\frac{d^2 x_i}{dt^2} = \sum_j \frac{\partial X_i}{\partial x_j} \frac{dx_j}{dt}, \quad i, j = 1, 2$$

or

$$(7) \quad \frac{d^2 x_1}{dt^2} = -\frac{dx_2}{dt}, \quad \frac{d^2 x_2}{dt^2} = \frac{dx_1}{dt}.$$

This prolongation admits the general solution

$$x_1(t) = a_1 \cos t + a_2 \sin t + h$$

$$x_2(t) = a_1 \sin t - a_2 \cos t + k, \quad t \in R.$$

(family of circles).

The pendulum geometric dynamics is described by

$$\frac{d^2 x_i}{dt^2} = \frac{\partial f}{\partial x_i} + \sum_j \left(\frac{\partial X_i}{\partial x_j} - \frac{\partial X_j}{\partial x_i} \right) \frac{dx_j}{dt}, \quad i, j = 1, 2$$

or

$$(8) \quad \frac{d^2 x_1}{dt^2} = x_1 - 2 \frac{dx_2}{dt}, \quad \frac{d^2 x_2}{dt^2} = x_2 + 2 \frac{dx_1}{dt},$$

with the general solution

$$x_1(t) = b_1 \cos t + b_2 \sin t + b_3 t \cos t + b_4 t \sin t$$

$$x_2(t) = b_1 \sin t - b_2 \cos t + b_3 t \sin t - b_4 t \cos t, \quad t \in R$$

(family of spirals).

Using

$$\begin{aligned}
L &= \frac{1}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 \right] + x_2 \frac{dx_1}{dt} - x_1 \frac{dx_2}{dt} + f \\
H &= \frac{1}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 \right] - f \\
g_{ij} &= (H + f) \delta_{ij}, \\
N_j^i &= -F_j^i = -\delta^{ih} F_{jh}, \quad F_{ij} = \frac{\partial X_j}{\partial x_i} - \frac{\partial X_i}{\partial x_j}, \quad i, j, h = 1, 2,
\end{aligned}$$

the solutions of the differential system (8) are horizontal pregeodesics of the Riemann-Jacobi-Lagrange manifold

$$(R^2 \setminus \{0\}, g_{ij}, N_j^i).$$

Lorenz Geometric Dynamics. We use the Riemannian manifold (R^3, δ_{ij}) . The Lorenz flow is a first dissipative model with chaotic behaviour discovered in numerical experiment. Its state equations are

$$\begin{aligned}
(9) \quad \frac{dx_1}{dt} &= -\sigma x_1 + \sigma x_2 \\
\frac{dx_2}{dt} &= -x_1 x_3 + r x_1 - x_2 \\
\frac{dx_3}{dt} &= x_1 x_2 - b x_3,
\end{aligned}$$

where σ, r, b are real parameters. Usually σ, b are kept fixed whereas r is varied. At

$$r > r_0 = \frac{\sigma(\sigma + b + 3)}{\sigma - b - 1}$$

chaotic behaviour is observed. With $\sigma = 10$, $b = \frac{8}{3}$, the preceding inequality yields $r_0 = 24,7368$. If $\sigma \neq 0$ and $b(r - 1) > 0$, then the equilibrium points of the Lorenz flow are

$$\begin{aligned}
&x = 0, \quad y = 0, \quad z = 0; \\
&x = \pm \sqrt{b(r - 1)}, \quad y = \pm \sqrt{b(r - 1)}, \quad z = r - 1.
\end{aligned}$$

Let

$$\begin{aligned}
X &= (X_1, X_2, X_3), \quad X_1(x_1, x_2, x_3) = -\sigma x_1 + \sigma x_2, \\
X_2(x_1, x_2, x_3) &= -x_1 x_3 + r x_1 - x_2, \quad X_3(x_1, x_2, x_3) = x_1 x_2 - b x_3, \\
f &= \frac{1}{2} [(-\sigma x_1 + \sigma x_2)^2 + (-x_1 x_3 + r x_1 - x_2)^2 + (x_1 x_2 - b x_3)^2], \\
rot X &= (2x_1, -x_2, r - x_3 - \sigma).
\end{aligned}$$

The Lorenz geometric dynamics is described by

$$\frac{d^2 x_i}{dt^2} = \frac{\partial f}{\partial x_i} + \sum_j \left(\frac{\partial X_i}{\partial x_j} - \frac{\partial X_j}{\partial x_i} \right) \frac{dx_j}{dt}, \quad i, j = 1, 2, 3$$

or

$$\begin{aligned}
(10) \quad \frac{d^2 x_1}{dt^2} &= \frac{\partial f}{\partial x_1} + (\sigma + x_3 - r) \frac{dx_2}{dt} - x_2 \frac{dx_3}{dt} \\
\frac{d^2 x_2}{dt^2} &= \frac{\partial f}{\partial x_2} + (r - x_3 - \sigma) \frac{dx_1}{dt} - 2x_1 \frac{dx_3}{dt} \\
\frac{d^2 x_3}{dt^2} &= \frac{\partial f}{\partial x_3} + x_2 \frac{dx_1}{dt} + 2x_1 \frac{dx_2}{dt}.
\end{aligned}$$

Using

$$\begin{aligned}
L &= \frac{1}{2} \sum_{i=1}^3 \left(\frac{dx_i}{dt} \right)^2 - \sum_{i=1}^3 X_i \frac{dx_i}{dt} + f \\
H &= \frac{1}{2} \sum_{i=1}^3 \left(\frac{dx_i}{dt} \right)^2 - f \\
g_{ij} &= (H + f) \delta_{ij} \\
N_j^i &= -F_j^i = -\delta^{ih} F_{jh}, \quad F_{ij} = \frac{\partial X_j}{\partial x_i} - \frac{\partial X_i}{\partial x_j}, \quad i, j, h = 1, 2, 3,
\end{aligned}$$

the solutions of the differential system (10) are horizontal pregeodesics of the Riemann-Jacobi-Lagrange manifold

$$(R^3 \setminus \mathcal{E}, \mathcal{J}|, \mathcal{N}^{\mathcal{J}}),$$

where $\mathcal{E}$ is the set of equilibrium points.

ABC Geometric Dynamics. We use the Riemannian manifold (R^3, δ_{ij}) . One examples of a fluid velocity that contains exponential stretching and hence instability is the ABC flow,

$$(11) \quad \begin{cases} \frac{dx_1}{dt} = A \sin x_3 + C \cos x_2 \\ \frac{dx_2}{dt} = B \sin x_1 + A \cos x_3 \\ \frac{dx_3}{dt} = C \sin x_2 + B \cos x_1. \end{cases}$$

This flow is named after the three mathematicians Arnold, Beltrami and Childress, who have contributed much to our understanding and appreciation of classes of "chaotic" flows of which the present one is an example. For nonzero values of the constants A, B, C the preceding system is not globally integrable. The topology of the flow lines is very complicated and can only be investigated numerically to reveal regions of chaotic behaviour. The ABC flow conserves the volumes since the ABC field is solenoidal.

The ABC geometric dynamics is described by

$$\frac{d^2 x^i}{dt^2} = \frac{\partial f}{\partial x^i} + \sum_j \left(\frac{\partial X_i}{\partial x_j} - \frac{\partial X_j}{\partial x_i} \right) \frac{dx_j}{dt}, \quad i, j = 1, 2, 3.$$

Since

$$f = \frac{1}{2}(A + B + C + 2AC \sin x_3 \cos x_2 + 2BA \sin x_1 \cos x_3 + 2CB \sin x_2 \cos x_1)$$

$$rot X = X,$$

the ABC geometric dynamics is given by the differential system (12):

$$\begin{aligned} \frac{d^2 x_1}{dt^2} &= AB \cos x_1 \cos x_3 - BC \sin x_1 \sin x_2 - (B \cos x_1 + C \sin x_2) \frac{dx_2}{dt} + \\ &\quad + (B \sin x_1 + A \cos x_3) \frac{dx_3}{dt} \\ \frac{d^2 x_2}{dt^2} &= -AC \sin x_2 \sin x_3 + BC \cos x_1 \cos x_2 + (B \cos x_1 + C \sin x_2) \frac{dx_1}{dt} - \\ &\quad - (A \sin x_3 + C \cos x_2) \frac{dx_3}{dt} \end{aligned}$$

$$\begin{aligned}\frac{d^2x_3}{dt^2} &= AC \cos x_3 \cos x_2 - BA \sin x_1 \sin x_3 - (B \sin x_1 + A \cos x_3) \frac{dx_1}{dt} + \\ &+ (C \cos x_2 + A \sin x_3) \frac{dx_2}{dt}.\end{aligned}$$

Using

$$\begin{aligned}L &= \frac{1}{2} \sum_{i=1}^3 \left(\frac{dx_i}{dt} \right)^2 - \sum_{i=1}^3 X_i \frac{dx_i}{dt} + f \\ H &= \frac{1}{2} \sum_{i=1}^3 \left(\frac{dx_i}{dt} \right)^2 - f \\ g_{ij} &= (H + f) \delta_{ij} \\ N_j^i &= -F_j^i = -\delta^{ih} F_{jh}, \quad F_{ij} = \frac{\partial X_j}{\partial x_i} - \frac{\partial X_i}{\partial x_j}, \quad i, j, h = 1, 2, 3\end{aligned}$$

the solutions of the differential system (12) are horizontal pregeodesics of the Riemann-Jacobi-Lagrange manifold

$$(R^3 \setminus \mathcal{E}, \}_{\rangle|}, \mathcal{N}_{\langle}^{\rangle}),$$

where $\mathcal{E}$ is the set of equilibrium points which is included in the surface of equation

$$\sin x_1 \sin x_2 \sin x_3 + \cos x_1 \cos x_2 \cos x_3 = 0.$$

References

- [1] R.G.Beil, *Comparison of unified field theories*, Tensor N.S., 56, 2(1995), 175-183.
- [2] R.G.Beil, *Notes on a new Finsler metric function*, Balkan Journal of Geometry and Its Applications, 2, 1(1997), 1-6.
- [3] F.M.Crampin, *A linear connection associated with any second order differential equation field*, In: Eds. L.Tamassy, J.Szenthe, New Developments in Geometry, 77-86, Kluwer Academic Publishers, 1996.
- [4] S.Friedlander, V.Yudovich, *Instabilities in fluid motions*, Notices of the AMS, 46, 11(1999), 1358-1367.

- [5] V.Obădeanu, C.Vernic, *Systemes dynamiques sur des espace de Riemann*, Balkan Journal of Geometry and Its Applications, 2 1(1997), 73-82.
- [6] A.Udrişte, C.Udrişte, *Dynamics induced by a magnetic field*, Conference on Differential Geometry, Budapest, July 27-30, 1996. In: Ed.J.Szenthe, New Developments in Differential Geometry, Kluwer Academic Publishers (1996), 429-442.
- [7] C.Udrişte, A.Udrişte, *Electromagnetic dynamical systems*, Balkan Journal of Geometry and Its Applications, 2,1(1997), 129-140.
- [8] C.Udrişte, *Geometric dynamics*, Second Conference of Balkan Society of Geometers, Aristotle University of Thessaloniki, June 23-26, 1998; Southeast Asian Bulletin of Mathematics 24, 1(2000), 1-11.
- [9] C.Udrişte, *Geometric dynamics*, Kluwer Academic Publishers, 2000.
- [10] C.Udrişte, *Nonclassical Lagrangian Dynamics and Potential Maps*, Proceedings of the Conference on Mathematics in Honour of Professor Radu Roşca at the Occasion of his Ninetieth Birthday, Katholieke University Brussel, Katolieke University Leuven, Belgium, Dec. 11-16, 1999.
- [11] C.Udrişte, *Dynamics induced by second-order objects*, BSG Proceedings 4, Global Analysis, Differential Geometry, Lie Algebras, Editor Grigorios Tsagas, pp.161-168, Geometry Balkan Press, 2000.
- [12] C.Udrişte, *Solutions of DEs and PDEs as potential maps using first order Lagrangians*, Centennial Vranceanu, Romanian Academy, University of Bucharest, June 30 - July 4, 2000.
- [13] C.Udrişte, M.Neagu, *Geometrical interpretation of solutions of certain PDEs*, Balkan Journal of Geometry and Its Applications, 4, 1(1999), 138-145.